

PCT/EP 04/13534



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03027383.3

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr:
Application no.: 03027383.3
Demande no:

Anmeldetag:
Date of filing: 27.11.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

DeveloGen Aktiengesellschaft für
entwicklungsbiologische Forschung
Rudolf-Wissell-Strasse 28
37079 Göttingen
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Method for preventing and treating diabetes using neurturin

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A61K38/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

27. NOV. 2003

WEICKMANN & WEICKMANN

Patentanwälte
European Patent Attorneys · European Trademark Attorneys

EPO - Munich
33

27 Nov. 2003

DIPL.-ING. H. WEICKMANN (bis 31.1.01)
DIPL.-ING. F. A. WEICKMANN
DIPL.-CHEM. B. HUBER
DR.-ING. H. LISKA
DIPL.-PHYS. DR. J. PRECHTEL
DIPL.-CHEM. DR. B. BÖHM
DIPL.-CHEM. DR. W. WEISS
DIPL.-PHYS. DR. J. TIESMEYER
DIPL.-PHYS. DR. M. HERZOG
DIPL.-PHYS. B. RUTTENSPERGER
DIPL.-PHYS. DR.-ING. V. JORDAN
DIPL.-CHEM. DR. M. DEY
DIPL.-FORSTW. DR. J. LACHNIT

Unser Zeichen:

31972P EP/WW/d

Anmelder:

DeveloGen Aktiengesellschaft für
entwicklungsbiologische Forschung
Rudolf-Wissell-Straße 28

37079 Göttingen
DEUTSCHLAND

Method for preventing and treating diabetes using neurturin

Method for preventing and treating diabetes using neurturin

Description

The present invention relates generally to methods for preventing and/or treating pancreatic disorders, particularly those related to diabetes, by
5 administering a neurturin product.

BACKGROUND OF THE INVENTION

The pancreas is an exocrine gland that secretes digestive enzymes directly
10 into the digestive tract as well as an endocrine gland that secretes hormones into the blood stream. The exocrine function is assured by acinar and centroacinar cells that secrete various digestive enzymes via intercalated ducts into the duodenum. The functional unit of the endocrine pancreas is the islet of Langerhans. Islets are scattered throughout the exocrine portion of the
15 pancreas and are composed of four main cell types: alpha-, beta-, delta- and PP-cells (reviewed for example in Kim & Hebrok, 2001, Genes Dev. 15: 111-127). Beta-cells produce insulin, represent the majority of the endocrine cells and form the core of the islets, while alpha-cells secrete glucagon and are located in the periphery. Delta-cells and PP-cells are less numerous and
20 secrete somatostatin and pancreatic polypeptide, respectively. Recently, cells producing the neuropeptide Ghrelin have been found in pancreatic islets (Wierup et al., 2002, Regul Pept. 107:63-9.).

Early pancreatic development has been well studied in different species,
25 including chicken, zebrafish, and mice (for a detailed review, see Kim & Hebrok, 2001, supra). The pancreas develops from distinct dorsal and ventral anlagen. Pancreas development requires specification of the pancreas anlage along both anterior-posterior and dorsal-ventral axes. A number of transcription factors, that are critical for proper pancreatic development have
30 been identified (see Kim & Hebrok, 2001, supra; Wilson et al., 2003, Mech Dev. 120: 65-80).

In humans, the acinar and ductal cells retain a significant proliferative capacity that can ensure cell renewal and growth, whereas the islet cells become mostly mitotically inactive. This is in contrast to rodents where beta-cell replication is an important mechanism in the generation of new beta cells. It has been suggested, that during embryonic development, pancreatic islets of Langerhans originate from differentiating duct cells or other cells with epithelial morphology (Bonner-Weir & Sharma, 2002, J Pathol. 197: 519-526; Gu et al., 2003, Mech Dev. 120: 35-43). In adult humans, new beta-cells arise in the vicinity of ducts (Butler et al., 2003, Diabetes 52: 102-110; Bouwens & Pipeleers 1998, Diabetologia 41: 629-633). However, also an intra-islet location or an origin in the bone marrow has been suggested for precursor cells of adult beta-cells (Zulewski et al., 2001, Diabetes 50: 521-533; lanus et al., 2003, J Clin Invest. 111: 843-850). Pancreatic islet growth is dynamic and responds to changes in insulin demand, for example during pregnancy or due to changing body weight during childhood. In adults, there is a good correlation between body mass and islet mass (Yoon et al., 2003, J Clin Endocrinol Metab. 88: 2300-2308).

Pancreatic beta-cells secrete insulin in response to blood glucose levels. Insulin amongst other hormones plays a key role in the regulation of the fuel metabolism. Insulin leads to the storage of glycogen and triglycerides and to the synthesis of proteins. The entry of glucose into muscles and adipose cells is stimulated by insulin. In patients who suffer from diabetes mellitus type I or LADA (latent autoimmune diabetes in adults (Pozzilli & Di Mario, 2001, Diabetes Care. 8:1460-67) beta-cells are being destroyed due to autoimmune attack. The amount of insulin produced by the remaining pancreatic islet cells is too low, resulting in elevated blood glucose levels (hyperglycemia). In diabetes type II liver and muscle cells lose their ability to respond to normal blood insulin levels (insulin resistance). High blood glucose levels (and also high blood lipid levels) in turn lead to an impairment of beta-cell function and to an increase in beta-cell apoptosis. It is interesting to note that the rate of beta-cell neogenesis does not appear to change in type II diabetics (Butler

et al., 2003 supra), thus causing a reduction in total beta-cell mass over time. Eventually the application of exogenous insulin becomes necessary in type II diabetics.

5 Improving metabolic parameters such as blood sugar and blood lipid levels (e.g. through dietary changes, exercise, medication or combinations thereof) before beta-cell mass has fallen below a critical threshold leads to a relatively rapid restoration of beta-cell function. However, after such a treatment the pancreatic endocrine function would remain impaired due to
10 the only slightly increased regeneration rate.

In type I diabetics, where beta-cells are being destroyed by autoimmune attack, treatments have been devised which modulate the immune system and may be able to stop or strongly reduce islet destruction (Raz et al., 2001,
15 Lancet 358: 1749-1753; Chatenoud et al., 2003, Nat Rev Immunol. 3: 123-132; Homann et al., Immunity. 2002, 3:403-15). However, due to the relatively slow regeneration of human beta-cells such treatments can only be successful if they are combined with agents that can stimulate beta-cell regeneration.

20 Diabetes is a very disabling disease, because today's common anti-diabetic drugs do not control blood sugar levels well enough to completely prevent the occurrence of high and low blood sugar levels. Out of range blood sugar levels are toxic and cause long-term complications like for example renopathy, retinopathy, neuropathy and peripheral vascular disease. There is also a host
25 of related conditions, such as obesity, hypertension, heart disease and hyperlipidemia , for which persons with diabetes are substantially at risk.

Apart from the impaired quality of life for the patients, the treatment of diabetes and its long term complications presents an enormous financial burden to our
30 healthcare systems with rising tendency. Thus, for the treatment of, type I and type II diabetes as well as for latent autoimmune diabetes in adults (LADA) there is a strong need in the art to identify factors that induce regeneration of pancreatic insulin producing beta-cells. These factors could restore normal

function of the endocrine pancreas once its function is impaired or event could prevent the development or progression of diabetes type I, diabetes type II, or LADA.

5 In this invention, we disclose a novel and so far unknown use for the neurotrophic factor neurturin to stimulate the formation or regeneration of insulin producing beta-cells and thus, a use in the treatment and prevention of diabetes.

10 Neurotrophic factors are growth factors that regulate the survival and maintain the phenotypic differentiation of certain nerve and/or glial cell populations (Varon et al., 1978, Ann. Rev. Neuroscience 1: 327-361; Thoenen et al., Science, 229:238-242, 1985). Nerve growth factor (NGF) was the first neurotrophic factor to be identified and characterized (Levi-Montalcini et al., 1951, J. Exp. Zool. 116:321). The second member of this
15 family to be discovered was brain-derived neurotrophic factor (Leibrock et al., 1989, Nature 341:149-152).

20 Glial-derived neurotrophic factor (GDNF) – a neurotrophic factor structurally unrelated to NGF - was discovered during a search for factors crucial to the survival of midbrain dopaminergic neurons, which degenerate in Parkinson's disease (Lin et al., 1993, Science 260:1130-2). Sequence analysis revealed it to be a distant member of the superfamily of transforming growth factor 8 (TGF-beta) factors.

25 Another neurotrophic factor that is structurally closely related to GDNF and unrelated to NGF is neurturin (Kotzbauer et al., 1996, Nature 384: 467-470). Neurturin, GDNF and two other related factors (artemin and persephin) define a family of neurotrophic factors referred to as TGF-beta-related
30 neurotrophins. These neurotrophic factors promote the survival of various neurons including peripheral autonomic and sensory neurons as well as central motor and dopamine neurons, and have been proposed as therapeutic agents for neurodegenerative diseases (see review by

Takahashi, 2001, Cytokine Growth Factor Rev 12(4):361-73; see also, for example, US 6,090,778 and EP1005358B1, the disclosures of which are hereby incorporated by reference). TGF-beta-related neurotrophins signal through a unique two-receptor complex consisting of a glycosylphosphatidylinositol-linked cell surface molecule, the GDNF family receptor alpha (GFRalpha), and receptor protein tyrosine kinase, Ret.

Apart from the described functions in neuronal tissue GDNF/RET signalling is crucial for the differentiation of certain non-neuronal tissues. GFRalpha1 and the transmembrane tyrosine kinase Ret are both necessary receptor components for ureteric bud outgrowth and subsequent branching in the developing kidney (Cacalano et al., 1998, Neuron 21:53-62; Tang et al., 1998, J Cell Biol. 142(5):1337-45). Ret, GFRalpha-1 (the GDNF receptor), and GFRalpha-2 (the Neurturin receptor) are expressed by testicular germ cells, while GDNF and Neurturin are expressed by Sertoli cells. Both GDNF and Neurturin stimulate DNA synthesis in spermatogonia. Furthermore, GFRalpha, GFRalpha ligands and co-receptors are expressed in germ cell tumors and thus may act as paracrine factors in spermatogenesis (Viglietto et al., 2000, Int J Oncol. 16(4):689-94)

Only recently it was shown that the biology of GDNF signalling is much more complex than originally assumed. GDNF family ligands also signal through the neural cell adhesion molecule NCAM. In cells lacking Ret, GDNF binds with high affinity to the NCAM and GFRalpha1 complex (see review by Sariola & Saarma, 2003, J Cell Sci. 116(Pt 19):3855-62). Signalling via the c-met receptor kinases has also been demonstrated (see Popsueva et al., 2003, J Cell Biol. 161(1):119-29).

SUMMARY OF THE INVENTION

The present invention relates to new methods for stimulating and/or inducing the differentiation of progenitor cells, e.g. stem cells into insulin-producing cells or for promoting the protection, survival and/or regeneration of insulin

producing cells using a neurturin product and/or a modulator/effector thereof that influences, particularly increases the expression level or function of a neurturin protein product.

5 Thus, the present invention provides methods for treating patients suffering from a disease caused by, associated with, and/or accompanied by functionally impaired and/or reduced numbers of pancreatic islet cells, particularly insulin producing beta-cells, by administering a therapeutically effective amount of a neurturin product or a compound that influences the
10 neurturin expression level or function. Functional impairment or loss of pancreatic islet cells may be due to e.g. autoimmune attack such as in diabetes type I or LADA, and/or due to cell degeneration such as in progressed diabetes type II. The methods of the present invention may also be used to treat patients at risk to develop degeneration of insulin producing
15 beta-cells to prevent the start or progress of such process.

The neurturin product or the effector/modulator thereof may be administered e.g. as a pharmaceutical composition, via implantation of neurturin product expressing cells and/or via gene therapy.

20

Further, the invention relates to cell preparations comprising neurturin-treated insulin producing cells or neurturin expressing cells.

Numerous additional aspects and advantages of the invention will become
25 apparent to those skilled in the art upon consideration of the following description of the Figures and detailed description of the invention which describes presently preferred embodiments thereof.

BRIEF DESCRIPTION OF THE FIGURES

30 **Figure 1 shows the neurturin dependent induction of the differentiation of insulin producing cells.**

Mouse embryonic stem (ES) cells were differentiated as described

previously (patent application PCT/EP02/04362, published as WO 02/086107, which is incorporated herein by reference). At the end of the differentiation procedure, cells were harvested and total RNA was isolated. The abundance of insulin mRNA (Figure 1 A) and of beta-cell glucose transporter Glut2 mRNA (Figure 1 B) was determined using quantitative RT-PCR in an Applied Biosystems 7000 sequence detection device. Levels were normalized using 18S RNA as control and a cycle number of 36 as reference. The numbers on the vertical line refer to the abundance of the indicated transcripts relative to an abundance for which 36 cycles are necessary for detection. ,wt ES' refers to unmodified mouse R1 embryonic stem (ES) cells; ,Pax4 ES' refers to R1 mouse embryonic stem (ES) cells stably transfected with a CMV-Pax4 expression construct; ,insulin expression rel. to delta Ct36' refers to expression of insulin; ,Glut2 expression rel. to delta Ct36' refers to expression of beta-cell glucose transporter; ,ES' refers to mouse embryonic stem cells, as described in Example 1; ,basic protocol' refers to the differentiation protocol as described in Example 2, without any addition of neurturin; ,EB+NTN' refers to the differentiation protocol as described in Example 3, with the addition of neurturin to embryoid bodies).

DETAILED DESCRIPTION OF THE INVENTION

Before the present invention is described, it is understood that all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs.

In the present invention the term "beta-cell regeneration" refers to an at least partial restoration of normal beta-cell function by increasing the number of functional insulin secreting beta-cells and/or by restoring normal function in functionally impaired beta-cells.

As used herein, the term "neurturin product" includes neurturin protein products such as purified natural or synthetic neurturin and variants thereof. Variants include insertion, substitution and deletion variants and chemically modified derivatives. Variants also include recombinant proteins, for

example but not limited to hybrids of neurturin and other TGF-beta proteins (preferably from the GDNF-family). Also included are proteins or peptides substantially homologous to the human neurturin precursor protein having the amino acid sequence published as GenBank Accession Number NP_004549. The term "neurturin product" also includes polynucleotides (e.g. mRNA/DNA) encoding the above described neurturin protein product. The term "neurturin product" also includes neurturin homodimers or heterodimers of a neurturin protein product and another protein, wherein the other protein preferably belongs to the GDNF-family.

The term "biologically active" as used herein means that the neurturin product induces and/or stimulates the differentiation of insulin producing cells from progenitor, e.g. stem cells and/or promotes the protection, survival, or regeneration of insulin producing cells, e.g. beta-cells. The biological activity of neurturin products may be determined as described in the Examples of the present application.

The term "substantially homologous" as used herein means having a degree of homology to the biologically active human neurturin product resulting from the cleavage of the neurturin precursor having the amino acid sequence published as GenBank Accession Number NP_004549 or to the human neurturin precursor itself, that is preferably in excess of 70%, most preferably in excess of 80%, and even more preferably in excess of 90% or 95%. The degree of homology between the mouse and the human protein is about 91 %, and it is contemplated that preferred mammalian neurturin proteins will have a similarly high degree of homology. Also included are proteins which are hybrids between neurturin and another TGFbeta-protein, preferably another member of the GDNF-family which retain the stimulatory effect on islet cell formation found in Neurturin. The percentage of homology or identity between a neurturin product and the human neurturin protein or a precursor or a nucleic acid coding therefor may be determined according to standard procedures, e.g. by using the BLAST algorithm. Preferably, the percentage of homology or identity is calculated as the percentage of

nucleotide or amino acid residues found in the smaller of the two sequences that align with identical nucleotides or amino acid residues in the sequence being compared, when four gaps in a length of 100 nucleotides or amino acids may be introduced to assist in that alignment. Also included as substantially homologous is any neurturin protein product which may be isolated by virtue of cross-reactivity with antibodies to the neurturin protein product or whose genes may be isolated through hybridization with the gene or with segments of the gene encoding the neurturin protein product.

10 In connection with the present invention, the term "progenitor cells" relates to undifferentiated cells capable of being differentiated into insulin producing cells. The term particularly includes stem cells, i.e. undifferentiated or immature embryonic, adult, or somatic cells that can give rise to various specialized cell types. The term "stem cells" can include embryonic stem
15 cells (ES) and primordial germ cells (EG) cells of mammalian, e.g. human or animal origin. Isolation and culture of such cells is well known to those skilled in the art (see, for example, Thomson et al., 1998, Science 282:1145-1147; Shambloott et al., 1998, Proc. Natl. Acad. Sci. USA 95:13726-13731; US 6,090,622; US 5,914,268; WO 0027995; Notarianni et al., 1990, J. Reprod. Fert. 41:51-56; Vassilieva et al., 2000, Exp. Cell. Res. 258:361-
20 373). Adult or somatic stem cells have been identified in numerous different tissues such as intestine, muscle, bone marrow, liver, and brain. WO 03/023018 describes a novel method for isolating, culturing, and differentiating intestinal stem cells for therapeutic use. In the pancreas, several indications suggest that stem cells are also present within the adult
25 tissue (Gu & Sarvetnick, 1993, Development 118:33-46; Bouwens, 1998, Microsc. Res. Tech. 43:332-336; Bonner-Weir, 2000, J. Mol. Endocr. 24:297-302).

30 Embryonic stem cells can be isolated from the inner cell mass of pre-implantation embryos (ES cells) or from the primordial germ cells found in the genital ridges of post-implanted embryos (EG cells). When grown in special culture conditions such as spinner culture or hanging drops, both ES

and EG cells aggregate to form embryoid bodies (EB). EBs are composed of various cell types similar to those present during embryogenesis. When cultured in appropriate media, EB can be used to generate in vitro differentiated phenotypes, such as extraembryonic endoderm, hematopoietic cells, neurons, cardiomyocytes, skeletal muscle cells, and vascular cells. We have previously described a method that allows EB to efficiently differentiate into insulin-producing cells (as described in patent application PCT/EP02/04362, published as WO 02/086107 and by Blyszczuk et al., 2003, Proc Natl Acad Sci U S A. 100(3):998-1003, which are incorporated herein by reference).

The term 'cultivation medium' means a suitable medium capable of supporting growth and differentiation of stem cells. The term 'differentiation medium' means a suitable medium for inducing the differentiation of stem cells into insulin-producing cells. The term 'terminal differentiation medium' means a suitable medium for terminal differentiation of insulin-producing cells. Examples of preferred media are described in WO/023018, which are herein incorporated by reference.

In this invention, we disclose a novel and so far unknown use for the neurotrophic factor neurturin to stimulate and/or induce the formation or regeneration of insulin producing cells and thus, a use in the treatment and prevention of diseases going along with (e.g. caused by, associated with or accompanied by) impaired beta-cell function, for example but not limited to diabetes mellitus. More particularly, the diseases are diabetes type I, diabetes type II or LADA.

The present invention is based on the finding that neurturin stimulates the differentiation of insulin producing cells from stem cells in vitro. Thus, a therapeutically effective amount of neurturin product may be administered to promote the regeneration of pancreatic beta-cells or to promote the formation of insulin-producing cells from stem cells or progenitor cells in vitro or in vivo. The present invention further relates to applications in the medical

field that directly arise from the method of the invention. Additionally, the present invention relates to applications for the identification and characterization of compounds with therapeutic medical effects or toxicological effects that directly arise from the method of the invention.

5

According to this invention the neurturin product may be administered

- i) as a pharmaceutical composition e.g. enterally, parenterally or topically, preferably directly to the pancreas,
 - ii) via implantation of neurturin protein product expressing cells, and/or
 - 10 iii) via gene therapy
- as described in more detail below.

Further, the neurturin expression level in a patient might be influenced by a neurturin modulator/effector administered

- 15 i) as a pharmaceutical composition e.g. enterally, parenterally or topically, preferably directly to the pancreas,
 - ii) via cell based therapy, and/or
 - iii) via gene therapy
- as described in more detail below.

20

The neurturin product or the neurturin modulator/effector, i.e. a pharmaceutically active substance influencing, particularly increasing the neurturin expression level or function may be administered in the above described manner alone or in combination with another pharmaceutical composition useful to treat beta-cell degeneration, for example hormones, growth factors or immune modulating agents.

25

A neurturin product or a modulator/effector thereof may be administered in patients suffering from a disease going along with impaired beta-cell function, for example but not limited to diabetes type I, LADA, or progressed diabetes type II. It is further contemplated that a neurturin product or the modulator/effector thereof may be administered preventively to patients at risk to develop beta-cell degeneration, like for example but not limited to

30

patients suffering from diabetes type II or LADA in early stages. A variety of pharmaceutical formulations and different delivery techniques are described in further detail below.

5 The present invention also relates to methods for differentiating progenitor cells into insulin-producing cells in vitro comprising

- (a) activating one or more pancreatic genes in a progenitor, e.g. stem cell (optional step, particularly if embryonic stem cells are used)
- (b) aggregating said cells to form embryoid bodies (optional step,
10 particularly if embryonic stem cells are used)
- (c) cultivating embryoid bodies or cultivating adult stem cells (e.g., duct cells) in specific differentiation media containing a neurturin protein product and/or a modulator/effector thereof under conditions wherein beta-cell differentiation is significantly enhanced, and
- 15 (d) identifying and selecting insulin-producing cells.

Activation of pancreatic genes may comprise transfection of a cell with pancreatic gene operatively linked to an expression control sequence, e.g. on a suitable transfection vector, as described in WO 03/023018, which is
20 herein incorporated by reference. Examples of preferred pancreatic genes are Pdx1, Pax4, Pax6, neurogenin 3 (ngn3), Nkx 6.1, Nkx 6.2, Nkx 2.2, HB 9, BETA2/Neuro D, Isl 1, HNF1-alpha, HNF1-beta and HNF3 of human or animal origin. Each gene can be used individually or in combination with at least one other gene. Pax4 is especially preferred.

25 Neurturin products, e.g. neurturin protein or nucleic acid products, are preferably produced via recombinant techniques because such methods are capable of achieving high amounts of protein at a great purity, but are not limited to products expressed in bacterial, plant, mammalian, or insect cell
30 systems.

Neurturin Protein Product

Recombinant neurturin protein product forms include glycosylated and non-glycosylated forms of the protein. In general, recombinant techniques involve isolating the genes encoding for neurturin protein product, cloning the gene
5 in suitable vectors and/or cell types, modifying the gene if necessary to encode a desired variant, and expressing the gene in order to produce the neurturin protein product.

Alternatively, a nucleotide sequence encoding the desired neurturin product
10 may be chemically synthesized. It is contemplated that a neurturin product may be expressed using nucleotide sequences that vary in codon usage due to the degeneration of the genetic code or allelic variations or alterations made to facilitate production of the protein product by the selected cell.

15 Kotzbauer et al., Nature 384:467-470, describe the identification of a mouse cDNA and amino acid sequence and a human cDNA and amino acid sequence for neurturin protein. The neurturin products according to this invention may be isolated or generated by a variety of means. Exemplary methods for producing neurturin products useful are described in patent
20 application WO 97/08196, the disclosures of which are hereby incorporated by reference. Also described are a variety of vectors, host cells, and culture growth conditions for the expression of neurturin protein, as well as methods to synthesize variants of neurturin protein product. Additional vectors suitable for the expression of neurturin protein product in E. coli are
25 disclosed in Patent No. EP 0 423 980, the disclosure of which is hereby incorporated by reference.

The molecular weight of purified neurturin indicates that in its biologically
30 active form the protein is a disulfide-bonded dimer. The material isolated after expression in a bacterial system is essentially biologically inactive, and exists as a monomer. Refolding is necessary to produce the biologically active disulfide-bonded dimer. Processes suitable for the refolding and maturation of the neurturin expressed in bacterial systems are substantially

similar to those described in WO93/06116. Standard in vitro assays for the determination of neurturin activity are also substantially similar to those determining GDNF activity as described in WO93/06116 and in U.S. Application Serial No. 08/535,681, and are hereby incorporated by reference.

Neurturin product variants are prepared by introducing appropriate nucleotide changes into the DNA encoding the polypeptide or by in vitro chemical synthesis of the desired polypeptide. It will be appreciated by those skilled in the art that many combinations of deletions, insertions, and substitutions can be made resulting in a protein product variant presenting neurturin biological activity.

Mutagenesis techniques for the replacement, insertion or deletion of one or more selected amino acid residues are well known to one skilled in the art (e.g., U.S. Patent No. 4,518,584, the disclosure of which is hereby incorporated by reference.)

Neurturin substitution variants have at least one amino acid residue of the human or mouse neurturin amino acid sequence removed and a different residue inserted in its place. Such substitution variants include allelic variants, which are characterized by naturally occurring nucleotide sequence changes in the species population that may or may not result in an amino acid change.

Chemically modified derivatives of neurturin protein products also may be prepared by one of skill in the art given the disclosures herein. The chemical moieties most suitable for derivatization include water soluble polymers. A water soluble polymer is desirable because the protein to which it is attached does not precipitate in an aqueous environment, such as a physiological environment. Preferably, the polymer will be pharmaceutically acceptable for the preparation of a therapeutic product or composition. One skilled in the art will be able to select the desired polymer based on such

considerations as whether the polymer/protein conjugate will be used therapeutically, and if so, the desired dosage, circulation time, resistance to proteolysis, and other considerations. A particularly preferred water-soluble polymer for use herein is polyethylene glycol. Attachment at residues important for receptor binding should be avoided if receptor binding is desired. One may specifically desire an N-terminal chemically modified protein.

The present invention contemplates use of derivatives which are prokaryote-expressed neurturin, or variants thereof, linked to at least one polyethylene glycol molecule, as well as use of neurturin, or variants thereof, attached to one or more polyethylene glycol molecules via an acyl or alkyl linkage. Pegylation may be carried out by any of the pegylation reactions known in the art. See, for example: Focus on Growth Factors, 3 (2):4-10, 1992; EP 0 154 316, the disclosure of which is hereby incorporated by reference; EP 0 401 384; and the other publications cited herein that relate to pegylation.

The present invention also discloses use of derivatives which are prokaryote-expressed neurturin, or variants thereof, linked to at least one hydrophobic residue, for example fatty acid molecule, as well as use of neurturin, or variants thereof, attached to one or more hydrophobic residues. For example, patent application published as WO 03/010185, which is hereby incorporated by reference, describes a method for producing acylated polypeptides in transformed host cells by expressing a precursor molecule of the desired polypeptide which are then to be acylated in a subsequent in vitro step.

Polynucleotides Encoding Neurturin Protein Product

The present invention further provides polynucleotides that encode neurturin protein products, whether recombinantly produced or naturally occurring.

A nucleic acid sequence encoding a neurturin protein product, can readily be obtained in a variety of ways, including, without limitation, chemical

synthesis, cDNA or genomic library screening, expression library screening, and/or PCR amplification of cDNA. These methods and others useful for isolating such nucleic acid sequences are set forth, for example, by Sambrook et al. (Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989), by Ausubel et al., eds (Current Protocols in Molecular Biology, Current Protocols Press, 1994), and by Berger and Kimmel (Methods in Enzymology: Guide to Molecular Cloning Techniques, vol. 152, Academic Press, Inc., San Diego, CA, 1987). Chemical synthesis of a nucleic acid sequence which encodes a neurturin protein product can also be accomplished using methods well known in the art, such as those set forth by Engels et al. (Angew. Chem. Intl. Ed., 28:716-734, 3 0 1989).

Included within the scope of this invention are neurturin product polynucleotides with the native signal sequence and other pre-pro sequences as well as polynucleotides wherein the native signal sequence is deleted and replaced with a heterologous signal sequence. The heterologous signal sequence selected should be one that is recognized and processed, i.e., cleaved by a signal peptidase, by the host cell. For prokaryotic host cells that do not recognize and process the native neurturin signal sequence, the signal sequence is substituted by a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, or heat-stable enterotoxin 11 leaders. For yeast secretion, the native neurturin signal sequence may be substituted by the yeast invertase, alpha factor, or acid phosphatase leaders. In mammalian cell expression the native signal sequence is satisfactory, although other mammalian signal sequences may be suitable.

Expression and cloning vectors generally include a nucleic acid sequence that enables the vector to replicate in one or more selected host cells.

Neurturin Product Pharmaceutical Compositions

Neurturin product pharmaceutical compositions typically include a

therapeutically effective amount of a neurturin product in admixture with one or more pharmaceutically and physiologically acceptable formulation. In addition to the active ingredients, neurturin product pharmaceutical compositions may contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries, which facilitate processing of the active compounds into preparations, which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, Pa.), the disclosure of which is hereby incorporated by reference.

Once the therapeutic composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, or dehydrated or lyophilized powder. Such formulations may be stored either in a ready to use form or in a form, e.g., lyophilized, requiring reconstitution prior to administration. The optimal pharmaceutical formulations will be determined by one skilled in the art depending upon considerations such as the route of administration and desired dosage. Such formulations may influence the physical state, stability, rate of in vivo release, and rate of in vivo clearance of the present neurturin proteins, variants and derivatives. Other effective administration forms, such as slow-release formulations, inhalant mists, or orally active formulations are also envisioned.

For example, in a sustained release formulation, the neurturin product may be bound to or incorporated into particulate preparations of polymeric compounds (such as polylactic acid, polyglycolic acid, etc.) or liposomes.

Administration/Delivery of Neurturin Product

The neurturin product may be administered by any suitable means, preferably enterally or parenterally or topically directly to the pancreas, as known to those skilled in the art. The specific dose may be calculated according to considerations of body weight, body surface area or organ size. Further refinement of the calculations necessary to determine the appropriate dosage for treatment involving each of the above mentioned

formulations is routinely made by those of ordinary skill in the art and is within the ambit of tasks routinely performed. Appropriate dosages may be ascertained through use of the established assays for determining dosages utilized in conjunction with appropriate dose-response data. The final
5 dosage regimen involved in a method for treating the above described conditions will be determined by the attending physician, considering various factors which modify the action of drugs, e.g., the age, condition, body weight, sex and diet of the patient, the severity of any infection, time of administration and other clinical factors. As studies are conducted, further
10 information will emerge regarding the appropriate dosage levels for the treatment of various diseases and conditions.

It is envisioned that the continuous administration or sustained delivery of a neurturin product may be advantageous for a given treatment. While
15 continuous administration may be accomplished via a mechanical means, such as with an infusion pump, it is contemplated that other modes of continuous or near continuous administration may be practiced. For example, chemical derivatization or encapsulation may result in sustained release forms of the protein having the effect of continuous presence, in
20 predictable amounts, based on a determined dosage regimen. Thus, neurturin protein products include proteins derivatized or otherwise formulated to effectuate such continuous administration.

Neurturin product cell therapy, i.e. pancreatic implantation of cells producing
25 neurturin protein product, is also contemplated. This embodiment would involve implanting cells capable of synthesizing and secreting a biologically active form of neurturin protein product into patients. Such neurturin protein product-producing cells may be cells that are natural producers of neurturin protein product or may be cells that are modified to express the protein.
30 Such modified cells include recombinant cells whose ability to produce a neurturin protein product has been augmented by transformation with a gene encoding the desired neurturin protein product in a vector suitable for promoting its expression and secretion. In order to minimize a potential

immunological reaction in patients being administered neurturin protein product of a foreign species, it is preferred that the cells producing neurturin protein product be of human origin and produce human neurturin protein product. Likewise, it is preferred that the recombinant cells producing neurturin protein product be transformed with an expression vector containing a gene encoding a human neurturin protein product. Implanted cells may be encapsulated to avoid infiltration of surrounding tissue. Human or nonhuman animal cells may be implanted in patients in biocompatible, semipermeable polymeric enclosures or membranes that allow release of neurturin protein product, but that prevent destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissue.

Alternatively, neurturin protein product secreting cells may be introduced into a patient in need intraportally via a percutaneous transhepatic approach using local anaesthesia. Between 3000 and 100 000 equivalent differentiated insulin-producing cells per kilogram body weight are preferably administered. Such surgical techniques are well known in the art and can be applied without any undue experimentation, see Pyzdrowski et al, 1992, New England J. Medicine 327:220-226; Hering et al., Transplantation Proc. 26:570-571, 1993; Shapiro et al., New England J. Medicine 343:230-238, 2000.

In a further preferred embodiment, neurturin protein product can be delivered directly to progenitor, e.g. stem cells in order to stimulate the differentiation of insulin producing cells. For example, protein delivery can be achieved by polycationic liposomes (Sells et al. (1995) Biotechniques 19:72-76), Tat-mediated protein transduction (Fawell et al. (1993) Proc. Natl. Acad. Sci. USA 91:664-668) and by fusing a protein to the cell permeable motif derived from the PreS2-domain of the hepatitis-B virus (Oess and Hildt (2000) Gene Ther. 7:750-758). Preparation, production and purification of such proteins from bacteria, yeast or eukaryotic cells are well known by persons skilled in the art. In this embodiment of the invention, neurturin may

be added preferably at concentrations between 1 ng/ml and 500 ng/ml, more preferably between 10 and 100 ng/ml, e.g. at about 50 ng/ml.

Further, the invention relates to a cell preparation comprising differentiated
5 progenitor cells, e.g. stem cells exhibiting insulin production, particularly an
insulin-producing cell line obtainable by the method described above. The
insulin-producing cells may exhibit a stable or a transient expression of at
least one pancreatic gene involved in beta-cell differentiation. The cells are
preferably human cells that are derived from human stem cells. For
10 therapeutic applications the production of autologous human cells from adult
stem cells of a patient is especially preferred. However, the insulin producing
cells may also be derived from non-autologous cells. If necessary, undesired
immune reactions may be avoided by encapsulation, immunosuppression
and/or modulation or due to non-immunogenic properties of the cells.

15 The insulin producing cells of the invention preferably exhibit characteristics
that closely resemble naturally occurring beta-cells. Further, the cells of the
invention preferably are capable of a quick response to glucose. After
addition of 27.7 mM glucose, the insulin production is enhanced by a factor
20 of at least 2, preferably by a factor of at least 3. Further, the cells of the
invention are capable of normalizing blood glucose levels after
transplantation into mice.

The invention further encompasses functional pancreatic cells obtainable or
25 obtained by the method according to the invention. The cells are preferably
of mammalian, e.g. human origin. Preferably, said cells are pancreatic beta-
cells, e.g. mature pancreatic beta-cells or stem cells differentiated into
pancreatic beta-cells. Such pancreatic beta cells preferably secrete insulin in
response to glucose. Moreover, the present invention provides functional
30 pancreatic cell that express glucagon in response to glucose. A preparation
comprising the cells of the invention may additionally contain cells with
properties of other endocrine cell types such as alpha-cells, delta-cells
and/or PP-cells. These cells are preferably human cells.

The cell preparation of the invention is preferably a pharmaceutical composition comprising the cells together with pharmacologically acceptable carriers, diluents and/or adjuvants. The pharmaceutical composition is preferably used for the treatment or prevention of pancreatic diseases, e.g. diabetes.

According to the present invention, the functional insulin producing cells treated with neurturin may be transplanted preferably intrahepatic, directly into the pancreas of an individual in need, or by other methods. Alternatively, such cells may be enclosed into implantable capsules that can be introduced into the body of an individual, at any location, more preferably in the vicinity of the pancreas, or the bladder, or the liver, or under the skin. Methods of introducing cells into individuals are well known to those of skill in the art and include, but are not limited to, injection, intravenous or parenteral administration. Single, multiple, continuous or intermittent administration can be effected. The cells can be introduced into any of several different sites, including but not limited to the pancreas, the abdominal cavity, the kidney, the liver, the celiac artery, the portal vein or the spleen. The cells may also be deposited in the pancreas of the individual.

The methodology for the membrane encapsulation of living cells is familiar to those of ordinary skill in the art, and the preparation of the encapsulated cells and their implantation in patients may be accomplished without undue experimentation. See, e.g., U.S. Patent Numbers 4,892,538, 5,011,472, and 5,106,627, each of which is specifically incorporated herein by reference. A system for encapsulating living cells is described in PCT Application WO 91/10425 of Aebischer et al., specifically incorporated herein by reference. See also, PCP Application WO 91/10470 of Aebischer et al., Winn et al., Exper. Neurol., 113:322-329, 1991, Aebischer et al., Exper. Neurol., 111:269-275, 1991; Tresco et al., ASAIO, 38:17-23, 1992, each of which is specifically incorporated herein by reference. Techniques for formulating a variety of other sustained- or controlled-delivery means, such as liposome carriers, bio-erodible particles or beads and depot injections, are also known

to those skilled in the art.

In another embodiment gene therapy ex vivo is envisioned, i.e. the patient's own cells may be transformed ex vivo to produce a neurturin protein product or a protein stimulating neurturin expression and would be directly reimplanted. For example, cells retrieved from the patient may be cultured and transformed with an appropriate vector. After an optional propagation/expansion phase, the cells can be transplanted back into the same patient's body, particularly the pancreas, where they would produce and release the desired neurturin protein product. Delivery by transfection and by liposome injections may be achieved using methods, which are well known in the art. Any of the therapeutic methods described above may be applied to any suitable subject including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

Neurturin product gene therapy in vivo is also envisioned, by introducing the gene coding for a neurturin protein product into targeted pancreas cells via local injection of a nucleic acid construct or other appropriate delivery methods (Hefti, J. Neurobiol., 25:1418-1435, 1994). For example, a nucleic acid sequence encoding a neurturin protein product may be contained in an adeno-associated virus vector or adenovirus vector for delivery to the pancreas cells. Alternative viral vectors include, but are not limited to, retrovirus, herpes simplex virus and papilloma virus vectors. Physical transfer, either in vivo or ex vivo as appropriate, may also be achieved by liposome-mediated transfer, direct injection (naked DNA), receptor-mediated transfer (ligand-DNA complex), electroporation, calcium phosphate precipitation or microparticle bombardment (gene gun).

Immunosuppressive drugs, such as cyclosporin, can also be administered to the patient in need to reduce the host reaction versus graft. Allografts using the cells obtained by the methods of the present invention are also useful because a single healthy donor could supply enough cells to regenerate at least partial pancreas function in multiple recipients.

Administration of a neurturin protein product and/or modulators/effectors thereof in a pharmaceutical composition to a subject in need thereof, particularly a human patient, leads to an at least partial regeneration of pancreatic cells. Preferably, these cells are insulin producing beta-cells that will contribute to the improvement of a diabetic state. With the administration of this composition e.g. on a short term or regular basis, an increase in beta-cell mass can be achieved. This effect upon the body reverses the condition of diabetes partially or completely. As the subject's blood glucose homeostasis improves, the dosage administered may be reduced in strength. In at least some cases further administration can be discontinued entirely and the subject continues to produce a normal amount of insulin without further treatment. The subject is thereby not only treated but could be cured entirely of a diabetic condition. However, even moderate improvements in beta-cell mass can lead to a reduced requirement for exogenous insulin, improved glycemic control and a subsequent reduction in diabetic complications. In another example, the compositions of the present invention will also have efficacy for treatment of patients with other pancreatic diseases such as pancreatic cancer, dysplasia, or pancreatitis, if beta-cells are to be regenerated.

In a further embodiment, the present invention allows the production of cells for the identification and/or characterisation of compounds which stimulate beta-cell differentiation, insulin secretion and/or glucose response, more particularly of compounds which increase the neurturin expression level or function. This method is particularly suitable for in vivo testing for diagnostic applications and drug development or screening. The compound of interest is added to suitable cells and the neurturin expression or function is determined. Alternatively, a compound of interest is added to a neurturin-treated cell and the effect on cell differentiation and/or insulin production is determined. Preferably, differentiated insulin-producing cells used. Insulin levels in treated cells can be determined, e.g. quantified by Enzyme Linked Immunoabsorbent Assay (ELISA) or Radio Immuno Assay (RIA). Using this method, a large number of compounds can be screened and compounds that induce neurturin expression or support the activity of neurturin leading

to a beta-cell differentiation and/or an increase in insulin secretion can be identified readily.

5 In a high-throughput screening method, the cells are transfected with a DNA construct, e.g. a viral or non-viral vector containing a reporter gene, e.g. the lacZ gene or the GFP gene, under regulatory control of a promoter of a gene involved in beta-cell differentiation, e.g. preferably a Pax4 promoter. The transfected cells are divided into aliquots and each aliquot is contacted with a test substance, e.g. candidate 1, candidate 2, and candidate 3. The
10 activity of the reporter gene corresponds to the capability of the test compound to induce beta-cell differentiation.

15 In a further embodiment (which may be combined with the high-throughput screening as described above) a medium throughput validation is carried out. Therein, the test compound is added to cells being cultivated and the neuritin expression and/or the insulin production is determined. Following an initial high throughput assay, such as the cell based assay outlined above where e.g. a Pax4 promoter is used as marker for beta-cell
20 differentiation is tested in a validation assay comprising adding said compounds to the culture media of the embryoid bodies. Differentiation into insulin-producing cells is then evaluated, e.g. by comparison to wild type and/or Pax4 expressing cells to assess the effectiveness of a compound.

25 **EXAMPLES**

A better understanding of the present invention and of its many advantages will be had from the following examples, given by way of illustration.

Example 1: Generation of ES cells expressing the Pax4 gene.

30 Mouse R1 ES cells (Nagy et al. (1993) Proc. Natl. Acad. Sci. U S A. 90:8424-8) were electroporated with the Pax4 gene under the control of the CMV promoter and the neomycin resistance gene under the control of the

phosphoglycerate kinase I promoter (pGK-1).

ES cells were cultured in Dulbecco's modified Eagle's medium containing 4.5 g/l glucose, 10^{-4} M beta-mercaptoethanol, 2 nM glutamine, 1% non-essential amino acids, 1 nM Na-pyruvate, 15% FCS and 500 U/ml leukaemia inhibitory factor (LIF). Briefly, approximately 10^7 ES cells resuspended in 0.8 ml phosphate buffered saline (PBS) were subjected to electroporation with 25 μ g/ml of linearized expression vector (Joyner, Gene Targeting: A Practical Approach, Oxford University Press, New York, 1993). Five minutes after electroporation, ES cells were plated on petri dishes containing fibroblastic feeder cells previously inactivated by treatment with 100 μ g/ml mitomycin C. One day after electroporation, culture medium was changed to medium containing 450 μ g/ml G418. Resistant clones were separately isolated and cultured 14 days after applying the selection medium. Cells were always cultured at 37°C, 5% CO₂. These untreated and undifferentiated ES cells were used as control the experiment shown in Figure 1 (referred to as 'ES' in Figure 1).

Example 2: Differentiation of ES cells into insulin-producing cells (referred to as 'basic protocol' in Figure 1)

The ES cell line R1 (wild type, 'wt ES' in Figure 1) and ES cells constitutively expressing Pax4 ('Pax4 ES' in Figure 1) were cultivated as embryoid bodies (EB) by the hanging drop method, as described in patent application PCT/EP02/04362, published as WO 02/086107 and by Blyszczuk et al., 2003, Proc Natl Acad Sci U S A. 100(3):998-1003, which are incorporated herein by reference. The embryoid bodies were allowed to form in hanging drop cultures for 2 days and then transferred for three days to suspension cultures in petri dishes. At day 5, EBs were plated separately onto gelatin-coated 24-well plates containing a differentiation medium prepared with a base of Iscove modified Dulbecco's medium. Cells were cultured for 15 to 40 days in the differentiation medium. To enhance differentiation capacity, a terminal differentiation medium can be applied at stages between 5 and 20

days after EB plating.

Example 3: Expression of pancreas specific genes after differentiation of ES cells into insulin-producing cells

5 Expression levels of pancreas specific genes was measured by semi-quantitative RT-PCR analysis. Differentiated wild type ES and Pax4 ES cells were collected after embryoid body formation and suspended in lysis buffer (4 M guanidinium thiocyanate, 25 mM sodium citrate, pH 7; 0.5% sarcosyl,
10 0.1 M beta-mercaptoethanol). Total RNA was isolated by the single step extraction method described by Chomczynski & Sacchi, 1987, Anal. Biochem. 162: 156-159). mRNA was reverse transcribed using PolyT tail primer Oligo d(T)₁₈ (PerkinElmer) and the resulting cDNA was amplified using oligonucleotide primers complementary and identical to transcripts of
15 beta-cell glucose transporter Glut2 and Insulin. The house keeping gene beta-tubulin was used as internal standard. Reverse transcription (RT) was performed with MuLV reverse transcriptase (Perkin Elmer). Multiplex PCRs were carried out using AmpliTaq DNA polymerase (Perkin Elmer) as described in Wobus et al., 1997, supra. mRNA levels of genes encoding
20 Glut2 and insulin were analysed using the Dynalbeads mRNA DIRECT micro kit (Dynal) according to the manufacturer's instructions.

One third of each PCR reaction was separated by electrophoresis. Ethidium bromide fluorescence signals of gels were analyzed by a special software
25 (TINA2.08e) The intensity of the ethidium bromide fluorescence signals was determined from the area under the curve for each peak and the data of target genes were plotted as percentage changes in relation to the expression of the housekeeping gene beta-tubulin.

30 Results show that markers for beta-cell differentiation function were expressed at higher levels in Pax4⁺ differentiated ES cells than in differentiated wild type ES cells demonstrating that activation of a pancreatic developmental control gene renders differentiation more efficient than for

wild type ES cells (Figure 1). Expression of Glut2 in differentiated stem cells indicates that hormone-producing cells are capable of responding to glucose. Expression of substantial amounts of insulin in differentiated stem cells indicates that differentiated cells show a phenotype similar to beta-cells.

Example 4: Induction of differentiation of insulin-producing cells by Neurturin (referred to as EB+NTN in Figure 1)

In order to study the effect of neurturin to induce beta-cell differentiation in vitro, we have generated stable mouse embryonic stem (ES) cells expressing the Pax4 under the control of the cytomegalovirus (CMV) early promoter/enhancer region as described in Example 1. Pax4 and wild type ES cells were then cultured in hanging drops or spinner cultures to allow the formation of embryoid bodies. Embryoid bodies were formed in the presence of 50 ng/ml of neurturin solution in PBS-01% BSA. Embryoid bodies were subsequently plated, again neurturin was added every second day until day 12 and afterwards cultured in a differentiation medium containing various growth factors (see Table 1 for more detail). Neurturin was obtained from RDI Research Diagnostics INC, USA, Order Number RDI-4511. Under such conditions, the expression of insulin was significantly induced by neurturin (Figure 1A). In addition, the addition of neurturin to the differentiation medium did significantly enhance the expression of the glucose transporter Glut-2 (Figure 1B). By comparison, wild type ES cells did not contain any insulin-producing cell at the same stage and only small numbers of glut-2 expressing cells. These data demonstrate that neurturin can significantly promote and enhance ES cells differentiation into insulin-producing cells compared to wild type ES cells.

The results shown in FIGURE 1 clearly demonstrate a significant induction of the differentiation of insulin-producing (Figure 1 A) and glucose responsive (Figure 1 B) cells, if neurturin is added to embryoid bodies. Thus, neurturin has a strong inductive effect on the differentiation of insulin-

producing beta cells.

Table 1. Protocol for the induction of differentiation of insulin-producing cells by Neurturin

5 Media B1, B2 and B27 are described in Rolletschek et al., 2001, Mech.Dev.;105, 2001:93-104.

Day	stage of cultivation	Medium	coating and analysis
0	Hanging Drops (200 cells/drop)	Iscove + 15% FCS Zugabe von Neurturin (NTN) 50 ng/ml	RNA (ES cells)
1			
2	EBs in Suspension	Iscove + 15%FCS + NTN, 50 ng/ml	
3			
4	Plating EBs	Iscove + 10%FCS + NTN, 50 ng/ml	Gelatin coating RNA (EBs)
+1	Change Medium	ITSFn medium + NTN, 50 ng/ml	Ornithin / Laminin coating
+2			
+3	"		
+4			
+5	"		
+6			
+7	"		
+8	Dissociation	N2+B27+NA+10%FCS	RNA (1x6 cm dish)
+9	Change Medium	s.o. withdraw FCS	
+10			
+11	"		
+12			
+13	"		
+14			
+15	"		
+16			
+17	"		
+18			
+19	"		
+20			
+21	"		
+22			
+23	"		
+24			
+25	"		Immunofluorescence (IF); RNA
+26			
+27	"		
+28			IF; RNA
+29	"		
+30			
+31	"		
+32			IF; RNA

Example 5: Functional characterisation of the differentiated insulin-producing cells

One important property of beta-cells is glucose responsive insulin secretion.

5 To test whether the Pax4 derived insulin-producing cells possessed this glucose responsive property, an *in vitro* glucose responsive assay was performed on the differentiated cells. Briefly, between 10 and 14 embryoid bodies were cultured in 3 cm petri dishes containing the above mentioned differentiation medium. On the day of the assay, the differentiation medium was removed and the cells are washed 3 times with Krebs Ringer Bicarbonate Hepes Buffer (KRBH; 118 mM NaCl, 4.7 KCl, 2.5 mM CaCl₂, 1,2 mM KH₂PO₄, 1.2 mM MgSO₄, 24.6 mM NaHCO₃, 10 mM Hepes, 2 mg/ml BSA). Cells were then incubated in 750 µl KRBH for 45 minutes at 37°C. The KRBH was then kept for measurement of basal insulin secretion and 750 µl 15 KRBH containing 27.7 mM glucose was added to the cells. After 15 minutes incubation at 37°C, the KRBH was removed from the cells for measurement of glucose-induced insulin secretion. Insulin levels were determined by an Enzyme-Linked Immunosorbent Assay (ELISA) for mouse insulin (Mercodia) and performed according to the manufacturer's recommendations. An 20 alternative medium for proper insulin release was medium based on DMEM with glucose concentration of 1 g/l (Gibco) supplemented with non-essential amino acids (Gibco, stock solution 1:100) and additional factors mentioned above. Such medium can be applied 1 to 6 days before use of the cells.

25 A basal insulin secretion is expected when both wild type and neurturin induced insulin-producing cells are cultured in the absence of glucose. However, only the neurturin induced insulin-producing cells respond to glucose stimulation. In the presence of glucose, a high increase in insulin secretion is expected in neurturin ES derived insulin-producing cells.

Example 6: Transplantation of Pax4 ES derived insulin-producing cells in STZ diabetic mice.

5 The therapeutic potential of neurturin induced insulin-producing cells to improve and cure diabetes can be investigated by transplanting the cells into streptozotocin induced diabetic mice. Streptozotocin is an antibiotic which is cytotoxic to beta-cells when administered at certain dosage (see Rodrigues et al.: Streptozotocin-induced diabetes, in McNeill (ed) Experimental Models of Diabetes, CRC Press LLC, 1999). Its effect is rapid, rendering an animal
10 severely diabetic within 48 hours.

Non-fasted Male BalbC mice were treated with STZ to develop hyperglycaemia after STZ treatment. Mice were considered diabetic if they had a blood glucose level above 10 mmol/l for more than 3 consecutive
15 days. Cells were transplanted under the kidney capsule and into the spleen of animals. The presence of the insulin-producing cells was confirmed by immunohistological analysis of the transplanted tissue. Results are expected to demonstrate that the transplanted cells can normalise blood glucose in diabetic animals.

20 All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific
25 preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are
30 intended to be within the scope of the following claims.

EPO - Munich
33
27 Nov. 2003

Claims

- 5 1. Use of a neurturin product and/or a modulator/effector thereof for the manufacture of a medicament to stimulate and/or induce the differentiation of insulin producing cells from progenitor cells.
2. The use of claim 1, wherein the progenitor cells are stem cells.
- 10 3. The use of claim 1, wherein the stem cells are embryonic or somatic stem cells.
4. The use of any one of claims 1-3, wherein the stem cells are of
15 mammalian origin, preferably of human origin.
5. The use of any one of claims 1-4, wherein the progenitor cells have been transfected with a pancreatic gene, particularly the Pax4 gene.
- 20 6. The use of a neurturin product and/or a modulator/effector thereof for the manufacture of a medicament to promote the protection, survival and/or regeneration of insulin producing cells.
7. The use of claim 6, wherein the insulin producing cells are beta-cells.
- 25 8. The use of claim 6 or 7, wherein the insulin producing cells are of mammalian origin, preferably of human origin.
9. The use of any one of claims 6-8, wherein the insulin producing cells
30 have been transfected with a pancreatic gene, particularly the Pax4 gene.

10. The use of any one of claims 1-9 for the prevention or treatment of a disease going along with impaired beta-cell function.

5 11. The use of claim 10 for the treatment of beta-cell degeneration in patients suffering from diabetes type I, LADA, or progressed diabetes type II.

10 12. The use of claim 10 for the prevention of beta-cell degeneration in patients at risk to develop beta-cell degeneration, like for example but not limited to patients suffering from diabetes type I or II, or LADA in early stages.

15 13. The use of any one of claims 1-12, wherein a neurturin product or a modulator/effector thereof that influences the expression level or function of a neurturin product is administered to a patient

- (i) as a pharmaceutical composition e.g. enterally, parenterally or topically directly to the pancreas,
- (ii) via implantation of neurturin protein product expressing cells, and/or
- 20 (iii) via gene therapy.

25 14. The use of claim 13, wherein the neurturin product or modulator/effector thereof is administered in combination with another pharmaceutical composition useful to treat beta-cell degeneration, for example but not limited to hormones, growth factors, or immune modulating agents.

30 15. The use of any one of claims 1-14, wherein the neurturin product is a protein including purified natural, synthetic or recombinant neurturin and variants thereof.

16. The use of claim 15 wherein variants are selected from insertion, substitution, deletion variants and/or chemically modified derivatives,

for example but not limited to hybrids of neurturin and other TGF-beta proteins preferably from the GDNF-family.

- 5 17. The use of claim 15 or 16, wherein the neurturin product is selected from proteins or peptides substantially homologous to the human neurturin precursor protein having the amino acid sequence published as GenBank Accession Number NP_004549 and/or to the mature neurturin protein product that results from the cleavage of the neurturin protein precursor published as GenBank Accession number
10 NP_004549.
18. The use of any one of claims 1-17, wherein the neurturin product is a nucleic acid, e.g. RNA and/or DNA encoding a neurturin protein product.
- 15 19. The use of any one of claims 1-18, wherein the neurturin product is selected from neurturin homodimers or heterodimers of a neurturin protein product and another protein, wherein the other protein preferably belongs to the GDNF-family or a nucleic acid coding therefor.
- 20 20. The use of any one of claims 1-19, wherein the neurturin product is of mammalian origin, preferably human origin.
- 25 21. The use of any one of claims 1-20, wherein the differentiation of progenitor, e.g. stem cells into insulin-producing cells in vitro comprises
- 30 a) optionally activating one or more pancreatic genes in progenitor cells,
b) optionally aggregating said cells to form embryoid bodies,
c) cultivating said cells or embryoid bodies in specific differentiation media containing neurturin protein product and
d) identifying and optionally selecting insulin-producing cells.

22. The use of claim 21, wherein the neurturin-treated insulin producing cells are

- (i) capable of a response to glucose and/or
- (ii) capable of expressing glucagon.

5

23. The use of any one of claims 21-22, wherein the neurturin-treated insulin producing cells are capable of normalizing blood glucose levels after transplantation into mice.

10

24. The use of any one of claims 1-23, wherein an effective amount of in vitro neurturin-treated cells are transplanted to a patient in need.

25. The use of any one of claims 1-24, comprising a stimulation of neurturin expression,

15

wherein cells from a patient in need that have been modified to produce and secrete a neurturin protein product in vitro are re-implanted into the patient and/or

wherein cells of a patient in need are modified to produce and secrete a neurturin protein product in vivo.

20

26. A method for differentiating or regenerating cells into functional pancreatic cells, the method comprising: (a) cultivating cells capable of being differentiated or regenerated into pancreatic cells in the presence of an effective amount of neurturin in vitro (b) allowing the cells to develop, to differentiate and/or to regenerate at least one pancreatic function; and (c) optionally preparing an effective amount of the differentiated or regenerated pancreatic cells for transplantation into a patient in need thereof.

25

30

27. The method of claim 26, wherein the patient in need is a human individual.

28. The method of claim 26 or 27, wherein the patient in need has (a)

functionally impaired, (b) reduced numbers and/or (c) functionally impaired and reduced numbers of pancreatic cells.

5 29. The method of any one of claims 26-28, wherein said patient in need is a type I diabetic patient or type II diabetic patient or LADA patient.

30. The method of any one of claims 26-29, wherein the pancreatic cells are insulin-producing cells.

10 31. The method of any one of claims 26-30, wherein the pancreatic cells are beta-cells of the pancreatic islets.

15 32. The method of any one of claims 26-31, wherein the cells in step (a) are selected from embryonic stem cells, adult stem cells, or somatic stem cells.

33. The method of any one of claims 27-32, wherein the cells in step (a) are of mammalian origin, preferably human origin.

20 34. The method of any one of claims 26-32, wherein neurturin is added at concentrations between 1 ng/ml and 500 ng/ml, preferably between 10 and 100 ng/ml, more preferably at about 50 ng/ml.

25 35. The method of any one of claims 26-34, wherein the at least one pancreatic function is selected from insulin production in response to glucose and expression of glucagon.

30 36. A method for differentiating or regenerating cells into functional pancreatic cells, the method comprising: preparing an effective amount of a neurturin product or of cells capable of expressing a neurturin product for administration to a patient in need thereof.

37. The method of claim 36, wherein the neurturin product is a neurturin protein product.

5 38. The method of of claim 36, wherein the neurturin product is a nucleic acid encoding a neurturin protein product.

39. The method of claim 36, wherein cells have been modified to produce and secrete a neurturin protein product and are prepared for transplantation into a suitable location in the patient.

10 40. A cell preparation comprising neurturin-treated functional pancreatic cells obtainable by the method of any one of claims 26-35.

15 41. A cell preparation comprising a neurturin product expressing cells obtainable by the method of any one of claims 36-40.

42. The preparation of claim 40 or 41, which is a pharmaceutical composition.

20 43. The preparation of any one of claims 40-42 for the treatment or prevention of pancreatic diseases, particularly diabetes.

44. The preparation of any one of claims 40-43 for administration by transplantation or for use in a medical device.

25 45. The preparation of any one of claims 40-44, which contains pharmaceutically acceptable carriers, diluents, and/or additives.

30 46. The preparation of any one of claims 40-45, which is a diagnostic composition.

47. The preparation of any one of claims 40-46, which is a therapeutic composition.

48. The preparation of any one of claims 40-47 for the manufacture of an agent for the regeneration of pancreatic tissues or cells, particularly pancreatic beta cells.

5

49. The preparation of any one of claims 40-48 for application in vivo.

50. The preparation of any one of claims 40-49 for application in vitro.

10 51. A method for identifying and/or characterizing compounds capable of modulating the differentiation or regeneration of cells into functional pancreatic, particularly insulin-producing cells comprising:
contacting a compound to be tested with cells under conditions wherein the cells are capable of being differentiated or regenerated into
15 functional pancreatic cells in the presence of neurturin and determining the effect of the compound on the differentiation process.

20 52. The method of claim 51 comprising transfecting the cells with a DNA construct containing a reporter gene under regulatory control of a gene involved in beta-cell differentiation, contacting said transfected cells with a compound to be tested and determining the activity of the reporter gene.

25 53. The method of claim 50 or 51 comprising contacting embryoid bodies which are cultivated in a differentiation medium enhancing beta-cell differentiation with a compound to be tested and determining differentiation into insulin-producing cells.

30 54. A method for identifying and/or characterizing compounds capable of modulating the differentiation or regeneration of cells into functional pancreatic, particularly insulin-producing cells comprising:
contacting a compound to be tested with cells under conditions wherein the cells are capable of being differentiated or regenerated into

functional pancreatic cells and determining the effect of the compound on the expression of neurturin.

- 5 55. Use of a preparation of neurturin expressing cells for the treatment and prevention of diabetes.
56. The use of claim 55 for inducing the regeneration of pancreatic cells.
- 10 57. The use of claim 56, wherein pancreatic cells are beta-cells of the islets.
58. Use of a preparation of neurturin-treated cells for the treatment and/or prevention of diabetes.
- 15 59. The use of claim 58 wherein the cells are differentiated progenitor cells capable of insulin production.

Abstract

5 The present invention relates generally to methods for preventing and/or
treating pancreatic disorders, particularly those related to diabetes, by
administering a neurturin protein product.

10

Id 27.11.2003

27. Nov. 2003




Figure 1

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP04/013534

International filing date: 29 November 2004 (29.11.2004)

Document type: Certified copy of priority document

Document details: Country/Office: EP
Number: 03027383.3
Filing date: 27 November 2003 (27.11.2003)

Date of receipt at the International Bureau: 11 February 2005 (11.02.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record.

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.